

Important Safety Information

For Rats, Mice, Ferrets, and Non-Human Primates:

Only administer Ethiq® XR by subcutaneous injection. Ethiq XR is not intended for intravenous, intra-arterial, intrathecal, intramuscular, or intra-peritoneal injection. Do not use in animals with pre-existing respiratory compromise.

Death has been reported when non-steroidal anti-inflammatory drugs (NSAIDs such as meloxicam and carprofen) and Ethiq XR have been administered concomitantly in mice.

Do not house rats on wood chip-type bedding after administration of Ethiq XR. **Pica involving wood chip type bedding can be lethal.**

Ethiq XR may cause sedation, decreased blood pressure, decreased heart rate, decreased gastrointestinal mobility, and respiratory depression. Use caution with concomitant administration of Ethiq XR with drugs that cause respiratory depression. Animals should be monitored for signs of decreased cardiovascular and respiratory function when receiving Ethiq XR.

The safety of Ethiq XR has not been evaluated in pregnant, lactating, neonatal, or immune-compromised animals.

For Humans:

Not for use in humans. Keep out of reach of children and pets.

Ethiq XR contains buprenorphine, a Schedule III controlled substance with an abuse potential similar to other Schedule III opioids, which may lead to overdose and death.

Ethiq XR should be handled appropriately to minimize the risk of misuse, abuse, addiction, and criminal diversion, including restriction of access, the use of accounting procedures, and proper disposal methods as appropriate to the laboratory setting and as required by law.

Ethiq XR should only be handled and administered by a veterinarian, veterinarian technician, or laboratory staff trained in the handling of potent opioids. Wear protective clothing when administering Ethiq XR to avoid direct contact with human skin, eyes, oral, or other mucus membranes which could result in absorption of buprenorphine and adverse reactions.

For more information, consult the Prescribing Information including the Boxed Warning.

BOXED WARNING**Abuse Potential**

ETHIQA XR contains buprenorphine, an opioid that exposes humans to risks of misuse, abuse, and addiction, which can lead to overdose and death. Use of buprenorphine may lead to physical dependence. The risk of abuse by humans should be considered when storing, administering, and disposing of ETHIQA XR. Persons at increased risk for opioid abuse include those with a personal or family history of substance abuse (including drugs or alcohol) or mental illness (e.g., depression).

Life-Threatening Respiratory Depression

Serious, life-threatening, or fatal respiratory depression may occur with accidental exposure to or with misuse or abuse of ETHIQA XR. Monitor for respiratory depression if human exposure to buprenorphine occurs. Misuse or abuse of buprenorphine by swallowing, snorting, or injecting poses a significant risk of overdose and death.

Accidental Exposure

Because of the potential for adverse reactions associated with accidental exposure, ETHIQA XR should only be administered by veterinarians, veterinary technicians, or laboratory staff who are trained in the handling of potent opioids. Accidental exposure to ETHIQA XR, especially in children, can result in a fatal overdose of buprenorphine.

Risks From Concurrent Misuse or Abuse with Benzodiazepines or Other CNS Depressants

Concurrent misuse or abuse of opioids with benzodiazepines or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death.

See HUMAN SAFETY WARNINGS for detailed information.